

LAKE MACQUARIE – WYONG REVIEW OF MONTHLY AMBIENT AIR QUALITY DATA APRIL 2016

NSW Environment Protection Authority

22 June 2016

Job Number 14030303B

Prepared by Todoroski Air Sciences Pty Ltd Suite 2B, 14 Glen Street Eastwood, NSW 2122 Phone: (02) 9874 2123 Fax: (02) 9874 2125 Email: info@airsciences.com.au

Lake Macquarie – Wyong

Review of Monthly Ambient Air Quality Data

April 2016

DISCLAIMER This report was prepared by Todoroski Air Sciences in good faith exercising all due care and attention, but no representation or warranty, express or implied, is made as to the relevance, accuracy, completeness or fitness for purpose of this document in respect of any particular user's circumstances. Users of this document should satisfy themselves concerning its application to, and where necessary seek expert advice in respect of, their situation. The views expressed within are not necessarily the views of the Office of Environment and Heritage (OEH) and may not represent OEH policy.

© Copyright State of NSW and the Office of Environment and Heritage

With the exception of photographs, the State of NSW and Office of Environment and Heritage are pleased to allow this material to be reproduced in whole or in part for educational and non-commercial use, provided the meaning is unchanged and its source, publisher and authorship are acknowledged. Specific permission is required for the reproduction of photographs.

TABLE OF CONTENTS

1	INTRODUCTION1						
2	PROJECT SCOPE1						
3	THE	PURPOSE OF AMBIENT MONITORING	.1				
	3.1	More about air quality	.2				
4	AIR	QUALITY MONITORING SITES	.3				
5	AIR	QUALITY CRITERIA AND STANDARDS	.4				
	5.1	Particulate matter	.4				
	5.1.1	PM _{2.5} concentrations	.4				
	5.2	Other air pollutants	. 5				
	5.3	Summary of applicable criteria for this review	. 5				
6	MET	EOROLOGICAL MONITORING DATA	.6				
7	AME	BIENT AIR QUALITY MONITORING DATA	. 8				
	7.1	Preamble	. 8				
	7.2	Analysis of Monitoring Data	. 8				
	7.3	PM ₁₀	. 8				
	7.4	PM _{2.5}	.9				
	7.5	NO ₂	.9				
	7.6	SO ₂					
	7.7	O ₃	10				
8	ANA	LYSIS OF ELEVATED POLLUTANT LEVELS	17				
9	CON	ICLUSIONS	18				
10	0 REFERENCES						

LIST OF TABLES

Table 4-1: Monitoring sites	3
Table 5-1: NSW EPA air quality impact assessment criteria for particulates	4
Table 5-2: NEPM Standards for ambient PM ₁₀ concentrations	4
Table 5-3: NEPM Standards for ambient PM _{2.5} concentrations	5
Table 5-4: NSW EPA air quality impact assessment criteria for air pollutants	5
Table 5-5: Air quality criteria and standards used in this review	6
Table 7-1: Maximum pollutant levels - April 2016	8

LIST OF FIGURES

Figure 4-1: Monitoring site locations	3
Figure 6-1: April windroses - Wallsend, Dora Creek, Marks Point, Wyee, Norah Head and Wyong	7
Figure 7-1: Lake Macquarie - Wyong 24-hour average PM ₁₀ levels – April 2016	11
Figure 7-2: Lake Macquarie - Wyong 24-hour average PM2.5 levels – April 2016	12
Figure 7-3: Lake Macquarie - Wyong 1-hour average NO ₂ levels – April 2016	13
Figure 7-4: Lake Macquarie - Wyong 1-hour average SO ₂ levels – April 2016	14
Figure 7-5: Lake Macquarie - Wyong 1-hour average O3 levels – April 2016	15
Figure 7-6: Lake Macquarie - Wyong rolling 4-hour average O ₃ levels – April 2016	16

LIST OF APPENDICIES

Appendix A – How to read a windrose


Appendix B – Monitoring Data (Graphical)

Appendix C – Monitoring Data (Tabulated)

EXECUTIVE SUMMARY

This report has been prepared by Todoroski Air Sciences for the NSW Environment Protection Authority (NSW EPA) and presents ambient air quality monitoring data recorded in the Lake Macquarie - Wyong region for the month of April 2016. The results indicate that the air quality was good in the Lake Macquarie - Wyong region during April.

The data summary (shown below) indicates that all data recorded in April were below the applicable short term criteria. Further details are provided in the report.



Lake Macquarie - Wyong Air Quality Pictorial Summary - April 2016

Lake Macquarie – Wyong Air Quality Tabular Summary - April 2016

Cit.	PM ₁₀ (μg/m ³)	PM _{2.5} (μg/m³) 4-hour average	SO₂ (μg/m³)	O ₃ (µg/m ³) 4-hour average ¹	Ο ₃ (μg/m ³)	NO2 (μg/m³) -hour averag	SO ₂ (μg/m³)
Site	2.	+-llour average		ality Impact Criteria		-nour averag	<u>se</u>
	50	25*	228	171	214	246	570
Wallsend	√	✓	✓	√	√	√	√
Wyong	√	✓	✓	√	√	√	√
Dora Creek	-	-	✓	-	-	√	√
Marks Point	-	-	√	-	-	√	1
Wyee	-	√	√	-	-	√	√
Wakefield HVAS	√	-	-	-	-	-	-
Teralba HVAS	✓	-	-	-	-	-	-
 ✓ - All data below applicable criteria 		 - Not appression 	plicable	* - Standard for PM _{2.}	5	¹ - Rollin	g average

 At least one elevated level above applicable criteria HVAS - High Volume Air Sampler concentrations (refer to Section 5.1)

1 INTRODUCTION

This report has been prepared by Todoroski Air Sciences on behalf of the NSW EPA. It provides a summary and analysis of the available ambient air quality and meteorological data collected in the Lake Macquarie - Wyong region during April 2016.

2 PROJECT SCOPE

The following outlines the scope of work for this project.

- Provide a monthly report written in plain English to the NSW EPA summarising and analysing available air quality data and meteorological information.
- The report will be published on the NSW EPA's website and will assess the available data from monitoring stations operated by the NSW Office of Environment and Heritage (OEH) at Wyong and Wallsend, and by industry at Wyee, Marks Point, Dora Creek, Wakefield and Teralba.
- The aim is to provide a simplified report that is accessible and contains results that would be clearly understood by the general public.

The work is for the period from September 2013 to June 2016.

3 THE PURPOSE OF AMBIENT MONITORING

It is important to note that the data presented in this report are from both NSW EPA and industry monitoring sites. The NSW EPA and the industry sites collect data for different purposes and this needs to be understood when comparing the data to the criteria.

NSW EPA monitoring sites are specifically designed to measure the likely levels of pollutants that the general population in the area would experience (i.e. an underlying population exposure level), whereas industry monitoring sites are specifically designed to measure maximum levels in a particular location that may be affected by a particular industry.

Data from NSW EPA sites can be compared with national air quality standards. Where the levels measured at NSW EPA monitoring sites are above the national standards on a prolonged and consistent basis, this indicates that some investigation of the potential cause of the issue may be warranted to determine whether any action on a regional level would reduce or better manage the pollutant levels. In the case of PM₁₀ and PM_{2.5}, it is noted that all data must be published, however days with exceptional events (e.g. bushfires and dust storms) may be excluded for the purpose of assessing compliance with the national standards.

Data from industry monitoring sites can be compared with NSW EPA impact assessment criteria. Where the levels measured at industry monitoring sites are above the applicable impact assessment criteria on a prolonged and consistent basis, this indicates that further investigation is warranted to determine the potential cause and what action is required by industry to reduce or better manage the pollutant.

Whether there is any harmful effect on an individual due to an air pollutant will depend on many additional factors, and not just on the measured level of a pollutant. These factors include the total exposure to the pollutant, individual circumstances (age, health, body mass, levels of pollutants at work), levels of other pollutants in the area, and many other factors.



Where pollutant levels are below the criteria generally, harm would not be expected to occur, but it does not follow that harm automatically occurs when pollutant levels are above the criteria.

The criteria serve to highlight potential issues with the levels of pollutants that may warrant more detailed examination. The criteria may also serve to prioritise action in various areas, for example areas with the highest pollutant levels and highest populations or highest exposure would be expected to receive priority action.

3.1 More about air quality

More information about air quality can be found via the following links:

- The Air Quality Index (AQI) was developed by the NSW EPA as an easily understood means of rating the pollutant level relative to its pollutant criteria.
 - o http://www.environment.nsw.gov.au/AQMS/aboutaqi.htm
- Aqicn.org provides near real-time AQI values for monitoring locations around the world. It should be noted that the AQI presented on this website is calculated differently to the NSW EPA AQI and is less stringent than those used in Australia, thus a direct comparison may not be valid.
 - http://aqicn.org/map/world/
- + The NSW OEH website air quality page provides hourly updates of the AQI and data readings from the NSW EPA monitoring sites, and can provide daily forecasts for Sydney and alerts for elevated levels at Wallsend and Wyong, for example. The web tool also presents near real-time wind and pollutant data readings overlaid on regional maps for the Upper Hunter and Newcastle.
 - o http://www.environment.nsw.gov.au/aqms/aqi.htm
- + The Lower Hunter Particle Characterisation Study aims to determine the composition of particulate samples collected at monitoring sites at Beresfield, Newcastle, Stockton and Mayfield, and to identify the potential major sources of fine particulates in Newcastle and the Lower Hunter. Progress reports are published on the OEH website provided below.
 - o http://www.environment.nsw.gov.au/aqms/lowhunterparticle.htm
- + The Air Emissions in My Community web tool presents the estimated emission quantities of various substances and their sources by postcode (and larger) sized areas in an easy to use graphical interface. This is one of the best inventories of emissions that is available, but it is important to appreciate that it cannot include all sources of emissions. It is important to also understand that pollutant emissions are not the same as the pollutant levels that this report presents. Emissions in a given area are one of several important factors that affect pollutant levels in an area, for example the dispersion of the emissions in the atmosphere and how the emissions are released are critical in determining the air quality pollutant levels.
 - o http://www.epa.nsw.gov.au/air/airemissionsapp/airemissionswebtool.aspx
- + The NSW Health website provides information on how air pollution affects health and steps for reducing your air pollution and limiting your exposure.
 - o http://www.health.nsw.gov.au/environment/air/Pages/default.aspx

4 AIR QUALITY MONITORING SITES

Figure 4-1 and **Table 4-1** summarise the locations and recorded parameters of the monitoring sites in the Lake Macquarie - Wyong region in April 2016.




Figure 4-1: Monitoring site locations

Table	1.1.	Monitorin	a citoc

Monitoring Station	Туре	Recorded Parameters	Recording Periods
Wallsend NSW EPA site		PM ₁₀ (TEOM), PM _{2.5} , NO ₂ , SO ₂ , O ₃ , WS, WD	Hourly/Daily
Wyong NSW EPA site		PM ₁₀ (TEOM), PM _{2.5} , NO ₂ , SO ₂ , O ₃ , WS, WD	Hourly/Daily
Marks Point Industry site		NO ₂ , SO ₂ , WS, WD	Hourly
Wyee Industry site		PM _{2.5} , NO ₂ , SO ₂ , WS, WD	Hourly
Dora Creek Industry site		NO ₂ , SO ₂ , WS, WD	Hourly
Norah Head BOM weather station		WS, WD	Hourly
Wakefield HVASIndustry siteTeralba HVASIndustry site		PM ₁₀ (HVAS)	Every 6th Day
		PM ₁₀ (HVAS)	Every 6th Day
PM ₁₀ - Particulate matter < 10	ım	NO ₂ - Nitrogen dioxide	WS - Wind speed
PM _{2.5} - Particulate matter < 2.5	μm	SO ₂ - Sulfur dioxide	WD - Wind direction
TEOM - Tapered Element Oscil	lating Microbalance	HVAS - High volume air sampler (which samples	BOM - Bureau of
(which samples air con	tinuously)	for a 24-hour period every 6 days)	Meteorology



5 AIR QUALITY CRITERIA AND STANDARDS

The sections below identify the key pollutants currently being monitored at the Lake Macquarie - Wyong air quality monitoring sites and the applicable air quality criteria and standards.

5.1 Particulate matter

Particulate matter consists of particles of varying size and composition. The total mass of all particles suspended in air is defined as the Total Suspended Particulate matter (TSP). The upper size range for TSP is nominally taken to be 30 micrometres (μ m) as in practice particles larger than 30 to 50 μ m will settle out of the atmosphere too quickly to be regarded as air pollutants.

The TSP is defined further into two sub-components. They are PM_{10} particles, particulate matter with aerodynamic diameters of 10µm or less, and $PM_{2.5}$, particulate matter with aerodynamic diameters of 2.5µm or less.

Table 5-1 summarises the air quality criteria that are relevant to particulate pollutants as outlined in the NSW EPA document *Approved Methods for the Modelling and Assessment of Air Pollutants in NSW* (**NSW DEC, 2005**).

Pollutant	Averaging Period	Criterion
Total suspended particulates (TSP)	Annual	90µg/m³
	Annual	30µg/m³
Particulate Matter < $10\mu m$ (PM ₁₀)	24-hour	50µg/m³

 Table 5-1: NSW EPA air quality impact assessment criteria for particulates

Source: NSW DEC, 2005

The National Environment Protection Council varied the Ambient Air Quality National Environment Protection Measure (NEPM) on 4 February 2016 (**NEPC, 2016**) through an amendment that added an annual average standard for PM_{10} concentrations. The updated NEPM removes the allowance for five exceedances of the 24-hour average PM_{10} concentration standard, and requires that all of the data are published, however days with exceptional events (e.g. bushfires and dust storms) may be excluded for the purpose of NEPM compliance assessment.

Table 5-2: NEPM Standards for ambient PM ₁₀ concentrations

Pollutant	Averaging Period	NEPM Standard
Particulate Matter < 10µm (PM _{2.5})	24-hour	50µg/m³
	Annual	25µg/m³

Source: NEPC, 2016

The NEPM standards apply to the average, or general exposure of a population, rather than to "hot spot" locations such as industry monitoring sites.

5.1.1 PM_{2.5} concentrations

The NSW EPA currently do not have impact assessment criteria for $PM_{2.5}$ concentrations, however standards for $PM_{2.5}$ concentrations are included in the recently updated NEPM (**NEPC, 2016**) which changed the $PM_{2.5}$ advisory reporting standards to national environment protection standards for $PM_{2.5}$, as shown in **Table 5-3**.

Table 5-3: NEPM Standards for ambient PM _{2.5} concentrations					
Pollutant	Averaging Period	NEPM Standard			
Particulate Matter < 2.5µm (PM _{2.5})	24-hour	25μg/m³			
	Annual	8μg/m³			

Source: NEPC, 2016

The NEPM standards apply to the average, or general exposure of a population, rather than to "hot spot" locations such as industry monitoring sites.

5.2 Other air pollutants

Nitrogen dioxide (NO₂) is reddish-brown in colour (at high concentrations) with a characteristic odour and can irritate the lungs and lower resistance to respiratory infections such as influenza. NO₂ belongs to a family of reactive gases called nitrogen oxides (NO_x). These gases form when fuel is burned at high temperatures, and mainly originate from motor vehicles, power generators and industrial boilers (USEPA, 2013). NO_x may also be generated by blasting activities. It is important to note that when formed, NO_2 is generally a small fraction of the total NO_x generated.

Sulfur dioxide (SO₂) is a colourless, toxic gas with a pungent and irritating smell. It commonly arises in industrial emissions due to the sulfur content of the fuel. SO₂ can have impacts upon human health and the habitability of the environment for flora and fauna. SO₂ emissions are a precursor to acid rain, which can be an issue in the northern hemisphere; however it is not known to be an issue in NSW.

Ozone (O_3) has a slight blue colour and is a reactive gas comprised of three oxygen atoms. It is typically found in the upper atmosphere, and forms what is referred to as the ozone layer which filters harmful ultraviolet radiation from the sun, and the near ground level in the troposphere. Tropospheric ozone forms through reactions between nitrogen oxides and volatile organic compounds (VOCs) in the presence of ultraviolet radiation. Tropospheric ozone is the main component of photochemical smog and can impact human health.

Table 5-4 summarises the air quality criteria that are relevant to NO₂, SO₂ and O₃ as outlined in the NSW EPA document Approved Methods for the Modelling and Assessment of Air Pollutants in NSW (NSW DEC, 2005).

Pollutant	Averaging period	Criterion
Nitrogon Diovido (NO.)	1-hour	246μg/m³
Nitrogen Dioxide (NO ₂)	Annual	62μg/m³
	10-minute	712μg/m ³
Sulfur Dioxide (SO ₂)	1-hour	570µg/m³
	24-hour	228µg/m³
	Annual	60μg/m³
	1-hour	214µg/m³
Ozone (O ₃)	Rolling 4-hour	171µg/m³

Table E. A. NSW/EDA air quality impact according to ritoria for air pollutants

Source: NSW DEC, 2005

5.3 Summary of applicable criteria for this review

The particulate and gaseous pollutants monitored in the Lake Macquarie – Wyong region have air quality criteria which are averaged over short and long time periods. Annually averaged criteria require a full year of data.



As this report only looks at one month of ambient air quality data, the annual average criteria are not applicable. The SO₂ 10-minute average criterion was not included as 10-minute monitoring data are not available. Therefore the criteria relevant to this assessment are those averaged over the shorter time periods (1-hour and 24-hours).

Table 5-5: Air quality criteria and standards used in this review					
Pollutant	Averaging Period	Туре	Concentration		
Particulate Matter < 10μm (PM ₁₀)	24-hour	Criterion/ Standard	50µg/m³		
Particulate Matter < 2.5µm (PM _{2.5})	24-hour	Standard	25μg/m³		
Nitrogen Dioxide (NO ₂)	1-hour	Criterion/ Standard	246µg/m³		
Sulfur Dioxide (SO ₂)	1-hour	Criterion/ Standard	570μg/m³		
Sullui Dioxide (SO ₂)	24-hour	Criterion/ Standard	228µg/m³		
$O_{\text{rens}}(O_{\lambda})$	1-hour	Criterion/ Standard	214µg/m³		
Ozone (O ₃)	Rolling 4-hour	Criterion/ Standard	171µg/m³		

Table 5-5 summarises the applicable air quality criteria and standards for this review.

METEOROLOGICAL MONITORING DATA 6

Representative wind speed and direction data have been obtained from the Lake Macquarie - Wyong air quality monitoring stations. The data are presented as a series of windroses. For an example of how to read a windrose, refer to Figure A-1 in Appendix A.

Figure 6-1 presents the April 2016 windroses for Wallsend, Dora Creek, Marks Point, Wyee, Norah Head and Wyong.

The figure shows that the meteorological stations recorded winds which varied depending on the local influence of environmental features such as terrain, vegetation and buildings. Overall the stations recorded varied wind directions with relatively high proportions of low and calm wind speeds.

The Norah Head weather station recorded wind speeds which were generally higher than those recorded at the other stations. This is expected as the Norah Head weather station is located in an unsheltered coastal location that would be largely influenced by sea breezes.




Figure 6-1: April windroses – Wallsend, Dora Creek, Marks Point, Wyee, Norah Head and Wyong

The meteorological stations recorded varied wind directions with relatively high proportions of low and calm wind speeds

7 AMBIENT AIR QUALITY MONITORING DATA

7.1 Preamble

The monitoring data in this report are presented in raw form as provided to Todoroski Air Sciences by the NSW EPA.

The 24-hour average data presented in this report have been averaged using the 1-hour average readings. Days which contain less than 75% data (less than 18 hours of 1-hour average data) have not been included in this report.

All of the monitoring data provided to Todoroski Air Sciences are presented in this report. The data are shown in the results and Appendices as relevant. 1-hour, 24-hour average and rolling annual average data are presented in a graphical format in **Appendix B** and 24-hour average data are presented in tabulated format in **Appendix C**.

7.2 Analysis of Monitoring Data

Table 7-1 presents a summary of the maximum pollutant levels measured during April 2016. The results indicate that pollutant levels recorded in April were below the applicable short term criteria at all monitors at all times.

Table 7-1: Maximum pollutant levels - April 2016							Γ		
	PM ₁₀	PM _{2.5}	SO ₂	O ₃	O ₃	NO ₂	SO ₂		
Site	(µg/m³)	(μg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)	(µg/m³)		
	24-hour	24-hour	24-hour	Rolling 4-	1-hour	1-hour	1-hour		
	average	average	average	hour average	average	average	average		
	Air Quality Impact Criteria								
	50	25*	228	171	214	246	570		
Wallsend	31.1	15.3	11.7	121.5	130.6	71.4	40.0		
Wyong	28.4	13.1	9.4	154.7	175.6	41.0	68.5		
Dora Creek	-	-	8.2	-	-	44.2	68.2		
Marks Point	-	-	6.0	-	-	39.2	86.2		
Wyee	-	18.9	7.1	-	-	47.0	35.0		
Wakefield	24.0	-	-	-	-	-	-		
HVAS	24.0								
Teralba HVAS	28.0	-	-	-	-	-	-		

Table 7-1: Maximum pollutant levels - April 2016

* Standard for PM_{2.5} concentrations (refer to Section 5.1)

- Not applicable

7.3 PM₁₀

Figure 7-1 presents all of the 24-hour average PM₁₀ monitoring results recorded in the Lake Macquarie - Wyong region in April 2016.

Relative to the Air Quality Index, as shown by the coloured bands in the figure, PM₁₀ levels were very good or good at all monitors at all times.

All data recorded at the Lake Macquarie - Wyong monitoring sites were less than the PM_{10} criterion level in April.

Figure B-1 to **Figure B-2** in **Appendix B** present the 1-hour average, 24-hour average and rolling annual average PM_{10} data in graphical form for each individual site. There is no criterion that applies to 1-hour average PM_{10} levels and these 1-hour results are not intended to be compared with the PM_{10}



criterion. It is a normal occurrence, and it is expected that in the normal environment 1-hour average PM_{10} levels will fluctuate more significantly than 24-hour average PM_{10} levels.

Figure B-1 to **Figure B-2** show the rolling annual average PM_{10} levels recorded at the Wallsend and Wyong monitors were $16.7\mu g/m^3$ and $15.2\mu g/m^3$ respectively at the end of April 2016. The rolling annual average levels can be compared to the annual average standard of $25\mu g/m^3$, however typically a calendar year of data are used to determine whether the annual average standard has been met.

7.4 PM_{2.5}

Figure 7-2 presents all of the 24-hour average PM_{2.5} monitoring data recorded in the Lake Macquarie - Wyong region in April 2016.

Relative to the Air Quality Index, as shown by the coloured bands in the figure, the data indicate that recorded PM_{2.5} levels were very good or good at all monitors at all times with the exception of the Wyee monitor which recorded two days with fair levels.

All data recorded at the Lake Macquarie - Wyong monitoring sites were below the 24-hour average $PM_{2.5}$ standard of $25\mu g/m^3$ in April 2016.

Figure B-3 to **Figure B-5** in **Appendix B** present the 1-hour average, 24-hour average and rolling annual average PM_{2.5} data in graphical form for each individual site. There is no criterion that applies to 1-hour average PM_{2.5} levels and these 1-hour results are not intended to be compared with the PM_{2.5} standard. It is a normal occurrence, and it is expected that in the normal environment 1-hour average PM_{2.5} levels will fluctuate more significantly than 24-hour average PM_{2.5} levels.

Figure B-3 to **Figure B-5** show the rolling annual average $PM_{2.5}$ levels recorded at the Wallsend, Wyong and Wyee monitors were $8.0\mu g/m^3$, $5.4\mu g/m^3$ and $9.4\mu g/m^3$ respectively at the end of April 2016. The rolling annual average levels can be compared to the annual average standard of $8\mu g/m^3$, however typically a calendar year of data are used to determine whether the annual average standard has been met.

It should be noted that the Wyee monitor has consistently recorded similar trends and generally higher PM_{2.5} levels than the Wallsend and Wyong monitors since June 2015. The Wyee PM_{2.5} monitor is currently under investigation by the operator to determine if the recorded levels are correct, i.e. whether any abnormally elevated level may be due to the calibration/ performance of the instrument or site factors.

7.5 NO₂

Figure 7-3 presents the 1-hour average NO₂ monitoring data recorded in the Lake Macquarie - Wyong region in April 2016.

Relative to the Air Quality Index, as shown by the coloured bands in the figure, the data indicate the NO_2 levels were very good at all monitors at all times.

All data were below the applicable criterion on all days.

9



7.6 SO₂

Figure 7-4 presents the 1-hour average SO₂ monitoring data recorded in the Lake Macquarie - Wyong region in April 2016.

Relative to the Air Quality Index, as shown by the coloured bands in the figure, the data indicate the SO₂ levels were very good at all monitors at all times.

All data were below the applicable criterion on all days.

7.7 O₃

Figure 7-5 presents the 1-hour average O₃ monitoring data recorded in the Lake Macquarie - Wyong region in April 2016.

Figure 7-6 presents the rolling 4-hour average O3 monitoring data recorded in the Lake Macquarie -Wyong region in April 2016.

Relative to the Air Quality Index, as shown by the coloured bands in the figures, the data indicate the 1-hour average levels of O₃ were generally very good or good. The Wallsend monitor recorded fair levels for three hours in April 2016.

The rolling 4-hour average levels were generally very good or good. The Wyong and Wallsend monitors recorded fair levels for three and four hours respectively.

All data were below the applicable criterion on all days.




Figure 7-1: Lake Macquarie - Wyong 24-hour average PM₁₀ levels – April 2016

The recorded PM₁₀ levels were very good or good at all monitors at all times. All data recorded at the Lake Macquarie - Wyong monitoring sites were less than the 24-hour average criterion of 50µg/m³.






Figure 7-2: Lake Macquarie - Wyong 24-hour average PM_{2.5} levels – April 2016

The recorded $PM_{2.5}$ levels were very good or good at all monitors at all times with the exception of the Wyee monitor which recorded two days with fair levels. All data recorded at the Lake Macquarie - Wyong monitoring sites were below the 24-hour average $PM_{2.5}$ standard of $25\mu g/m^3$.






Figure 7-3: Lake Macquarie - Wyong 1-hour average NO₂ levels – April 2016

All data recorded at the Lake Macquarie - Wyong monitoring sites were below the 1-hour average NO₂ criterion level of $246\mu g/m^3$ in April 2016. The measured levels of NO₂ were very good at all monitors at all times.






Figure 7-4: Lake Macquarie - Wyong 1-hour average SO₂ levels - April 2016

All data recorded at the Lake Macquarie - Wyong monitoring sites were below the 1-hour average SO_2 criterion level of $570\mu g/m^3$ in April 2016. The measured levels of SO_2 were very good at all monitors at all times.






Figure 7-5: Lake Macquarie - Wyong 1-hour average O₃ levels – April 2016

All data recorded at the Lake Macquarie - Wyong monitoring sites were below the 1-hour average O_3 criterion level of $214\mu g/m^3$ in April 2016. Measured 1-hour average levels of O_3 were generally very good or good. The Wallsend monitor recorded fair levels for three hours in April 2016.






Figure 7-6: Lake Macquarie - Wyong rolling 4-hour average O₃ levels – April 2016

All data recorded at the Lake Macquarie - Wyong monitoring sites were below the rolling 4-hour average O_3 criterion level of 171μ g/m³ in April 2016. The measured levels were generally very good or good. The Wyong and Wallsend monitors recorded fair levels for three and four hours respectively.



ANALYSIS OF ELEVATED POLLUTANT LEVELS 8

There were no levels above the applicable criteria in April 2016.

CONCLUSIONS

9

The results indicate that the monitoring stations recorded good air quality during April 2016.

Relative to the Air Quality Index:

- The measured levels of NO₂ were very good at all monitors at all times;
- ✤ The measured levels of SO₂ were very good at all monitors at all times;
- + The measured levels of O₃ were generally very good or good. The Wyong and Wallsend monitors recorded fair levels for three and four hours respectively;
- + The measured PM_{2.5} levels were very good or good at all monitors at all times with the exception of the Wyee monitor which recorded two days with fair levels; and,
- The measured PM₁₀ levels were very good or good at all monitors at all times. +

All recorded rolling annual average levels were less than or equal to the applicable annual (calendar year) standards in April 2016, except for the Wyee monitor which recorded a rolling annual average $PM_{2.5}$ level of 9.4µg/m³ at the end of April 2016.

It should be noted that the Wyee monitor has consistently recorded similar trends and generally higher PM_{2.5} levels than the Wallsend and Wyong monitors since June 2015. The Wyee PM_{2.5} monitor is currently under investigation by the operator to determine if the recorded levels are correct, i.e. whether any abnormally elevated level may be due to the calibration/ performance of the instrument or site factors.

On this basis it can be concluded that the air quality in the Lake Macquarie - Wyong region was good in April 2016.



10 REFERENCES

NEPC (2001)

"National Environment Protection (Ambient Air Quality) Measure Technical Paper No. 5 Data Collection and Handling", National Environment Protection Council, May 2001.

NEPC (2016)

"National Environment Protection (Ambient Air Quality) Measure", National Environment Protection Council, February 2016.

NSW DEC (2005)

"Approved Methods for the Modelling and Assessment of Air Pollutants in NSW", Department of Environment and Conservation (NSW), August 2005.

USEPA (2013)

Health Effects of Pollution, United States Environmental Protection Agency website.<http://www.epa.gov/region07/air/quality/health.htm>, accessed May 2013.





Appendix A

How to read a windrose






Figure A-1: How to read a windrose



Appendix B

Monitoring Data (Graphical)






Figure B-1: Wallsend PM₁₀ (1-hour, 24-hour and rolling annual average) concentration - April






Figure B-2: Wyong PM₁₀ (1-hour, 24-hour and rolling annual average) concentration – April






Figure B-3: Wallsend PM_{2.5} (1-hour, 24-hour and rolling annual average) concentration – April






Figure B-4: Wyong PM_{2.5} (1-hour, 24-hour and rolling annual average) concentration – April






Figure B-5: Wyee PM_{2.5} (1-hour, 24-hour and rolling annual average) concentration - April



Appendix C

Monitoring Data (Tabulated)



Date	ΡΜ ₁₀ (μg/m³)		Table C-1: April 24-hour avera PM _{2.5} (μg/m ³)			SO ₂ (μg/m ³)				
Date	Wallsend	Wyong	Wallsend	Wyong	Wyee	Wallsend	Wyong	Dora Creek	Marks Point	Wyee
1/04/2016	19.1	17.9	6.4	5.9	10.3	4.2	4.6	2.3	1.1	5.5
2/04/2016	23.1	24.9	10.5	10.7	14.3	1.0	0.4	1.6	1.7	0.8
3/04/2016	31.1	28.4	8.3	8.8	12.9	1.1	0.0	2.0	2.1	2.8
4/04/2016	18.6	21.0	6.0	5.6	9.9	4.6	2.5	5.4	1.3	1.0
5/04/2016	20.0	17.6	8.2	5.6	11.6	3.2	8.6	4.0	4.0	0.9
6/04/2016	27.4	23.2	12.4	9.0	12.7	3.5	1.4	2.9	3.7	1.2
7/04/2016	27.8	26.0	10.8	6.5	9.3	0.0	0.0	1.9	0.7	0.6
8/04/2016	17.2	14.5	3.5	5.2	8.3	0.5	0.1	4.8	1.1	0.2
9/04/2016	19.7	16.2	10.2	9.1	13.4	3.1	0.0	3.2	1.9	0.7
10/04/2016	24.5	21.2	12.6	13.1	17.2	6.1	0.2	3.3	3.6	0.8
11/04/2016	25.8	23.5	-	12.7	18.9	11.7	0.4	6.2	6.0	1.0
12/04/2016	18.4	16.0	15.3	9.1	12.4	0.6	0.2	5.5	1.0	2.4
13/04/2016	11.0	9.6	7.1	4.6	8.1	0.0	0.4	6.3	0.6	2.2
14/04/2016	11.8	7.5	4.7	4.0	6.2	2.5	0.0	1.8	1.3	3.0
15/04/2016	12.3	9.5	-	2.9	8.2	2.9	4.6	3.4	3.1	3.4
16/04/2016	14.7	13.0	9.9	6.6	8.5	3.1	4.0	2.2	1.9	0.7
17/04/2016	15.3	16.2	7.9	8.8	11.3	0.6	0.0	1.2	0.8	2.6
18/04/2016	14.0	12.9	6.3	6.4	9.5	3.6	0.5	1.0	1.7	-
19/04/2016	11.6	10.1	7.8	3.3	8.1	3.7	0.0	2.7	5.0	-
20/04/2016	12.7	8.8	6.6	3.4	9.8	0.5	0.0	3.2	0.9	3.9
21/04/2016	12.3	11.7	6.9	5.5	12.2	1.7	3.0	6.9	2.9	5.5
22/04/2016	20.5	18.5	10.9	9.3	15.1	3.5	0.4	3.0	0.9	6.9
23/04/2016	14.7	11.8	8.0	3.6	9.6	8.6	0.0	1.9	2.5	4.3
24/04/2016	23.0	19.8	8.9	4.9	10.0	0.7	0.0	7.3	0.7	4.8
25/04/2016	16.3	12.3	5.7	4.8	8.2	0.0	0.0	7.6	0.7	4.2
26/04/2016	11.5	10.7	6.0	4.3	8.3	0.9	0.0	2.3	0.8	3.9
27/04/2016	11.7	8.6	6.9	3.1	9.6	2.2	4.2	2.7	1.7	5.4
28/04/2016	11.2	22.1	-	4.1	10.2	-	9.4	8.2	2.4	5.7
29/04/2016	13.5	17.9	-	6.3	11.1	-	3.0	4.3	1.5	6.1
30/04/2016	14.8	12.5	7.9	5.4	11.2	4.4	1.6	3.2	5.8	7.1

Table C-1: April 24-hour average monitoring data

- Not applicable

Table C-2: April 24-hour average HVAS monitoring data

Data	PM ₁₀ (HVAS) (μg/m³)				
Date	Wakefield (Westside)	Teralba			
5/04/2016	9	22			
11/04/2016	24	28			
17/04/2016	18	16			
23/04/2016	15	20			
29/04/2016	11	24			

- Not applicable



C-1