



REVIEW OF NAMOI
AMBIENT AIR QUALITY DATA
AUTUMN 2017

NSW Environment Protection Authority

25 August 2017

Job Number 17050703

Prepared by

Todoroski Air Sciences Pty Ltd

Suite 2B, 14 Glen Street

Eastwood, NSW 2122

Phone: (02) 9874 2123

Fax: (02) 9874 2125

Email: info@airsciences.com.au

REVIEW OF NAMOI

AMBIENT AIR QUALITY DATA

AUTUMN 2017

DISCLAIMER This report was prepared by Todoroski Air Sciences in good faith exercising all due care and attention, but no representation or warranty, express or implied, is made as to the relevance, accuracy, completeness or fitness for purpose of this document in respect of any particular user's circumstances. Users of this document should satisfy themselves concerning its application to, and where necessary seek expert advice in respect of, their situation. The views expressed within are not necessarily the views of the Office of Environment and Heritage (OEH) and may not represent OEH policy.

© Copyright State of NSW and the Office of Environment and Heritage

With the exception of photographs, the State of NSW and Office of Environment and Heritage are pleased to allow this material to be reproduced in whole or in part for educational and non-commercial use, provided the meaning is unchanged and its source, publisher and authorship are acknowledged. Specific permission is required for the reproduction of photographs.

TABLE OF CONTENTS

1	INTRODUCTION.....	6
2	PROJECT SCOPE.....	6
3	THE PURPOSE OF AMBIENT MONITORING.....	6
3.1	More about air quality	6
4	AIR QUALITY MONITORING SITES	8
5	AIR QUALITY CRITERIA.....	9
5.1	Particulate matter.....	9
5.2	Summary of applicable criteria for this review.....	9
6	METEOROLOGICAL MONITORING DATA.....	10
7	AMBIENT AIR QUALITY MONITORING DATA	12
7.1	Preamble.....	12
7.2	Analysis of Monitoring Data	12
7.3	PM ₁₀	12
7.4	PM _{2.5}	13
8	ANALYSIS OF ELEVATED POLLUTANT LEVELS.....	17
8.1	Tamworth TEOM monitor – 10 April 2017.....	17
9	CONCLUSIONS.....	20
10	REFERENCES	21

LIST OF TABLES

Table 4-1: Monitoring sites	8
Table 5-1: NSW EPA air quality impact assessment criteria	9
Table 5-4: Air quality criteria used in this review	9
Table 7-1: Maximum and annual average pollutant levels – Autumn 2017	12

LIST OF FIGURES

Figure 4-1: Monitoring site locations	8
Figure 6-1: Autumn 2017 windroses – Tamworth, Werris Creek, Wil-gai, Gunnedah and Narrabri.....	11
Figure 7-1: Summary of AQI levels recorded in the Namoi region during autumn 2017	14
Figure 7-1: Namoi 24-hour average PM ₁₀ levels – Autumn 2017	15
Figure 7-2: Namoi 24-hour average PM _{2.5} levels – Autumn 2017	16
Figure 8-1: Analysis of elevated PM ₁₀ levels on 10 April 2017 - Tamworth.....	18
Figure 8-2: UHAQMN and Tamworth monitoring data on 10 April 2017	19

LIST OF APPENDICES


Appendix A – How to read a windrose	
Appendix B – Monitoring Data (Graphical)	
Appendix C – Monitoring Data (Tabulated)	

EXECUTIVE SUMMARY

This report has been prepared by Todoroski Air Sciences for the NSW Environment Protection Authority (NSW EPA). It presents ambient air quality monitoring data recorded in the Namoi region for the autumn 2017 period.

The results indicate that the air quality was generally very good in the Namoi region in autumn 2017. The data summary (shown below) indicates that the Tamworth monitor recorded one day with 24-hour average PM₁₀ levels above the criterion of 50µg/m³. All other data were below the applicable criteria. Further details are provided in the report. The 24-hour average data are provided in the Appendices.

Namoi Air Quality Pictorial Summary – Autumn 2017



Namoi Air Quality Tabular Summary – Autumn 2017

Site	PM ₁₀	PM _{2.5}
	Maximum 24-hour average	
	Air Quality Impact Criteria (µg/m ³)	
	50	25
Breeza	✓	✓
Maules Creek	✓	✓
Werris Creek	✓	✓
Wil-gai	✓	✓
Tamworth	✗	✓

✓ - All data below applicable criteria

✗ - At least one elevated level above applicable criteria

1 INTRODUCTION

This report has been prepared by Todoroski Air Sciences on behalf of the NSW EPA. It provides a summary and analysis of the available ambient air quality and meteorological data collected in the Namoi region in autumn 2017.

2 PROJECT SCOPE

The following outlines the scope of work for this project.

- ✦ Provide three reports for the Namoi region which cover the periods July 2015 to December 2016, summer (December 2016 to February 2017) and autumn (March 2017 to May 2017). The reports will examine compliance with 24-hour average and annual average criteria, summarise all of the reported data and include seasonal trends and analysis to identify likely source categories for elevated pollution events.
- ✦ The report will be published on the NSW EPA's website and will assess the available data from monitoring stations operated by the NSW Office of Environment and Heritage (OEH) at Tamworth, and by industry at Breeza, Maules Creek, Werris Creek and Wil-gai.
- ✦ The aim is to provide a simplified report that is accessible and contains results that would be clearly understood by the general public.

3 THE PURPOSE OF AMBIENT MONITORING

It is important to note that the data presented in this report are from both NSW EPA and industry monitoring sites. The NSW EPA and the industry monitoring sites collect data for different purposes and this needs to be understood when comparing the data with the criteria.

NSW EPA monitoring sites are specifically designed to measure the likely levels of pollutants that the general population in the area would experience (i.e. an underlying population exposure level), whereas industry monitoring sites are specifically designed to measure maximum levels in a particular location which may be affected by a particular industry.

Data from NSW EPA monitoring sites can generally be compared with national air quality standards. Where the levels measured at NSW EPA monitoring sites are above the national standards on a prolonged and consistent basis, this indicates that some investigation of the potential cause of the issue may be warranted to determine whether any action on a regional level would reduce or better manage the pollutant levels. In the case of PM₁₀ and PM_{2.5}, it is noted that all data must be published, however days with exceptional events (e.g. bushfires and dust storms) may be excluded for the purpose of assessing compliance with the national standards.

Data from industry monitoring sites can be compared with NSW EPA impact assessment criteria, and Project Approval criteria. Where the levels measured at industry monitoring sites are above the applicable impact assessment or Project Approval criteria on a prolonged and consistent basis, this indicates that further investigation is warranted to determine the potential cause and what action is required by industry to reduce or better manage the pollutant.

Whether there is any harmful effect on an individual due to an air pollutant will depend on many additional factors, and not just on the measured level of a pollutant. These factors include the total

exposure to the pollutant, individual circumstances (age, health, body mass, levels of pollutants at work), levels of other pollutants in the area, and many other factors. Where pollutant levels are below the criteria generally, harm would not be expected to occur, but it does not follow that harm automatically occurs when pollutant levels are above the criteria.

The criteria serve to highlight potential issues with the levels of pollutants that may warrant more detailed examination. The criteria may also serve to prioritise action in various areas, for example areas with the highest pollutant levels and highest populations or highest exposure would be expected to receive priority action.

3.1 More about air quality

More information about air quality can be found via the following links:

- + The NSW EPA website provides ambient air quality monitoring data on a weekly basis for four industry operated monitoring locations in the Namoi region, at Breeza, Maules Creek, Werris Creek and Wil-gai.
 - o <http://www.epa.nsw.gov.au/air/namoi/namoiairmon.htm>
- + The Air Quality Index (AQI) was developed by the NSW EPA as an easily understood means of rating the pollutant level relative to its pollutant criteria.
 - o <http://www.environment.nsw.gov.au/AQMS/aboutaqi.htm>
- + The NSW OEH website air quality page provides hourly updates of the AQI and data readings from the NSW EPA monitoring sites. Subscribers can sign up for alerts for elevated levels for the North-west slopes, based on the Tamworth monitoring site.
 - o <http://www.environment.nsw.gov.au/aqms/subscribe.htm>
- + Aqicn.org provides near real-time AQI values for monitoring locations around the world. It should be noted that the AQI presented on this website is calculated differently to the NSW EPA AQI and is less stringent than those used in Australia, thus a direct comparison may not be valid.
 - o <http://aqicn.org/map/world/>
- + The NSW Health website provides information on how air pollution affects health and steps for reducing your air pollution and limiting your exposure.
 - o <http://www.health.nsw.gov.au/environment/air/Pages/default.aspx>

4 AIR QUALITY MONITORING SITES

Figure 4-1 and Table 4-1 summarise the locations and recorded parameters of the monitoring sites in the Namoi region in autumn 2017.




Figure 4-1: Monitoring site locations

Table 4-1: Monitoring sites

Monitoring Station	Type	Recorded Parameters	Recording Periods
Tamworth	NSW OEH site	PM ₁₀ (TEOM), PM _{2.5} (BAM), WS, WD	Hourly/Daily
Breeza	Industry site	PM ₁₀ (TEOM), PM _{2.5} (TEOM), WS ¹ , WD ¹	Hourly/Daily
Maules Creek	Industry site	PM ₁₀ (TEOM), PM _{2.5} (TEOM), WS ¹ , WD ¹	Hourly/Daily
Werris Creek	Industry site	PM ₁₀ (TEOM), PM _{2.5} (TEOM), WS, WD	Hourly/Daily
Wil-gai	Industry site	PM ₁₀ (TEOM), PM _{2.5} (TEOM), WS, WD	Hourly/Daily
Gunnedah	BOM weather station	WS, WD	Hourly
Narrabri	BOM weather station	WS, WD	Hourly

PM₁₀ - Particulate matter < 10µm
 PM_{2.5} - Particulate matter < 2.5µm
 BOM - Bureau of Meteorology

TEOM - Tapered Element Oscillating Microbalance
 BAM - Beta Attenuation Monitor
¹ Sensor not at 10m above ground level

WS - Wind speed
 WD - Wind direction

5 AIR QUALITY CRITERIA

The sections below identify the key pollutants currently being monitored at the Namoi air quality monitoring sites and the applicable air quality criteria.

5.1 Particulate matter

Particulate matter consists of particles of varying size and composition. The total mass of all particles suspended in air is defined as the Total Suspended Particulate matter (TSP). The upper size range for TSP is nominally taken to be 30 micrometres (μm) as in practice particles larger than 30 to 50 μm will settle out of the atmosphere too quickly to be regarded as air pollutants.

The TSP is defined further into two sub-components. They are PM_{10} particles, particulate matter with aerodynamic diameters of 10 μm or less, and $\text{PM}_{2.5}$, particulate matter with aerodynamic diameters of 2.5 μm or less.

Table 5-1 summarises the air quality goals that are relevant to particulate pollutants as outlined in the NSW EPA document *Approved Methods for the Modelling and Assessment of Air Pollutants in New South Wales* (NSW EPA, 2017).

Table 5-1: NSW EPA air quality impact assessment criteria

Pollutant	Averaging Period	Criterion
Total suspended particulates (TSP)	Annual	90 $\mu\text{g}/\text{m}^3$
Particulate Matter < 10 μm (PM_{10})	Annual	25 $\mu\text{g}/\text{m}^3$
	24-hour	50 $\mu\text{g}/\text{m}^3$
Particulate Matter < 2.5 μm ($\text{PM}_{2.5}$)	Annual	8 $\mu\text{g}/\text{m}^3$
	24-hour	25 $\mu\text{g}/\text{m}^3$

Source: NSW EPA, 2017

5.2 Summary of applicable criteria for this review

The particulate pollutants monitored in the Namoi region have air quality criteria which are averaged over short and long time periods.

As this report looks at less than an annual period of ambient air quality data, only the criteria applicable averaged over the shorter time periods (24-hours) are applicable.

Table 5-2 summarises the applicable air quality criteria for this review.

Table 5-2: Air quality criteria used in this review

Pollutant	Averaging Period	Type	Concentration
Particulate Matter < 10 μm (PM_{10})	24-hour	Criterion / NEPM Standard ¹	50 $\mu\text{g}/\text{m}^3$
Particulate Matter < 2.5 μm ($\text{PM}_{2.5}$)	24-hour	Criterion/ NEPM Standard ¹	25 $\mu\text{g}/\text{m}^3$

¹ Source: NEPC, 2016

6 METEOROLOGICAL MONITORING DATA

Representative wind speed and direction data have been obtained from the relevant monitoring stations listed in **Table 4-1**.

For an example of how to read a windrose, refer to **Figure A-1** in **Appendix A**.

Figure 6-1 presents the 2017 autumn seasonal windroses for Tamworth, Werris Creek, Wil-gai, Gunnedah and Narrabri.

The windroses show that the meteorological stations recorded winds which varied over distance and depending on the local influence of environmental features such as terrain, vegetation and buildings.

The meteorological stations generally recorded winds from the southeast quadrant during autumn. The recorded wind speeds were generally higher at the Narrabri and Gunnedah (airport) weather stations than at the other sites. This is likely due to the clear open spaces and flat terrain which are characteristic of both sites. The Tamworth weather station generally recorded much lower wind speeds which is likely due to the siting of the station amongst buildings and vegetation within the township of Tamworth.




Figure 6-1: Autumn 2017 windroses – Tamworth, Werris Creek, Wil-gai, Gunnedah and Narrabri

The windroses show that the meteorological stations recorded winds which were generally from the southeast quadrant.

7 AMBIENT AIR QUALITY MONITORING DATA

7.1 Preamble

The monitoring data in this report are presented in raw form as provided to Todoroski Air Sciences by the NSW EPA.

The 24-hour average data presented in this report have been averaged using 1-hour average readings. Only data labelled as valid have been included in the averages. Days which contain less than 75% valid data (less than 18 hours of 1-hour average data) have not been included in this report.

All of the monitoring data provided to Todoroski Air Sciences are presented in this report. The data are shown in the results and Appendices as relevant. 1-hour, 24-hour and rolling annual average data are presented in a graphical format in **Appendix B** and 24-hour average data are presented in a tabulated format in **Appendix C**.

7.2 Analysis of Monitoring Data

Table 7-1 presents a summary of the pollutant levels measured during autumn 2017. The results indicate that the Tamworth monitor recorded a 24-hour average PM₁₀ level above the criterion of 50µg/m³. All other pollutant levels were below the applicable criteria.

Figure 7-1 presents a summary of the PM₁₀ and PM_{2.5} AQI levels recorded in the Namoi region during autumn 2017. The data indicate that the air quality was generally very good in the Namoi region during this period.

Table 7-1: Maximum and annual average pollutant levels – Autumn 2017

Site	PM ₁₀	PM _{2.5}
	Maximum 24-hour average	
	Air Quality Impact Criteria (µg/m ³)	
	50	25
Breeza	24.1	13.4
Maules Creek	34.7	6.9
Werris Creek	25.1	11.4
Wil-gai	41.8	11.3
Tamworth	54.1	14.9

7.3 PM₁₀

Figure 7-2 presents all of the 24-hour average PM₁₀ monitoring results recorded in the Namoi region during autumn 2017.

Relative to the Air Quality Index, as shown by the coloured bands in the figure, PM₁₀ levels were generally very good or good during autumn 2017. The Wil-gai monitor recorded fair levels on four days, and the Maules Creek and Tamworth monitors recorded one day with fair levels. The Tamworth monitor recorded one day with poor levels.

On 10 April 2017 the Tamworth monitor recorded elevated PM₁₀ levels above the 24-hour average criterion of 50µg/m³. All other 24-hour average data were below the criterion in autumn 2017.

Figure B-1 to **Figure B-5** in **Appendix B** present the 1-hour average, 24-hour average and rolling annual average PM₁₀ data in graphical form for each individual site. The rolling annual averages are

based on the previous 12-month periods of data. There is no criterion that applies to 1-hour average PM_{10} levels and these 1-hour results are not intended to be compared with the PM_{10} criterion. It is a normal occurrence, and it is expected that in the normal environment 1-hour average PM_{10} levels will fluctuate more significantly than 24-hour average PM_{10} levels.

We note the monitoring sites on occasion recorded periods in which PM_{10} levels were less than zero. In some situations the concentration of the pollutant being measured may be very close to zero, in which case the measured value may be less than the measurement limit of detection (**NEPC, 2001**), and in these circumstances the output may be negative.

The monitors may also record short term positive or negative values due to instrument faults, the presence of moisture within the instrument or volatile matter (which can register as a solid mass at first, but then evaporates, registering negative mass at a later time).

7.4 $PM_{2.5}$

Figure 7-3 presents all of the 24-hour average $PM_{2.5}$ monitoring data recorded in the Namoi region during autumn 2017.

Relative to the Air Quality Index, as shown by the coloured bands in the figure, the data indicate that $PM_{2.5}$ levels were generally very good in autumn 2017. The Maules Creek monitor recorded very good levels 100% of the time and the other monitors recorded very good levels between 65% (Breeza) and 93% (Wil-gai) of the time, with good levels recorded in remaining periods.

All 24-hour average data recorded at the Namoi monitoring sites were below the criterion in autumn 2017.

Figure B-6 to **Figure B-10** in **Appendix B** present the 1-hour average, 24-hour average and rolling annual average $PM_{2.5}$ data in graphical form for each individual site. The rolling annual averages are based on the previous 12-month periods of data. There is no criterion that applies to 1-hour average $PM_{2.5}$ levels and these 1-hour results are not intended to be compared with the $PM_{2.5}$ criteria. It is a normal occurrence, and it is expected that in the normal environment 1-hour average $PM_{2.5}$ levels will fluctuate more significantly than 24-hour average $PM_{2.5}$ levels.

We note the monitoring sites on occasion recorded periods in which $PM_{2.5}$ levels were less than zero. In some situations the concentration of the pollutant being measured may be very close to zero, in which case the measured value may be less than the measurement limit of detection (**NEPC, 2001**), and in these circumstances the output may be negative.

The monitors may also record short term positive or negative values due to instrument faults, the presence of moisture within the instrument or volatile matter (which can register as a solid mass at first, but then evaporates, registering negative mass at a later time).




Figure 7-1: Summary of AQI levels recorded in the Namoi region during autumn 2017

The data indicate that PM₁₀ levels in the Namoi region were very good between 67.6% (Wil-gai) and 97.8% (Werris Creek) of the time, and PM_{2.5} levels were very good between 64.7% (Breeza) and 100% (Maules Creek) of the time. PM_{2.5} levels were very good or good 100% of the time.




Figure 7-2: Namoi 24-hour average PM₁₀ levels – Autumn 2017

The recorded PM₁₀ levels were generally very good or good during autumn 2017. The Wil-gai monitor recorded fair levels on four days, and the Maules Creek and Tamworth monitors recorded one day with fair levels. The Tamworth monitor recorded one day with poor PM₁₀ levels above the 24-hour average criterion of 50µg/m³ which may have been caused by a regional dust storm. All other 24-hour average data were below the criterion in autumn 2017.




Figure 7-3: Namoi 24-hour average PM_{2.5} levels – Autumn 2017

The recorded PM_{2.5} levels were generally very good in autumn 2017. The Maules Creek monitor recorded very good levels 100% of the time and the other monitors recorded very good levels between 65% (Breeza) and 93% (Wil-gai) of the time, with good levels recorded in remaining periods. All data recorded at the Namoi monitoring sites were below the criterion of 25µg/m³.

8 ANALYSIS OF ELEVATED POLLUTANT LEVELS

There was one elevated level above the applicable criteria of $50\mu\text{g}/\text{m}^3$ in autumn 2017, this was;

- ✦ 24-hour average PM_{10} level of $50.9\mu\text{g}/\text{m}^3$ recorded at Tamworth on 10 April 2017.

8.1 Tamworth TEOM monitor – 10 April 2017

Figure 8-1 presents a plot of the 1-hour average PM_{10} , wind speed and wind direction data recorded at Tamworth on 10 April 2017, along with the PM_{10} levels recorded at the other Namoi monitors. The data presented in **Figure 8-1** show that the Tamworth monitor experienced elevated PM_{10} levels during a period of winds from the west and northwest from approximately 4am to 3pm. The other Namoi monitors recorded relatively high PM_{10} levels earlier in the morning and lower levels after midday.

Figure 8-2 presents the PM_{10} , wind speed and wind direction monitoring data from the Tamworth monitoring site (dashed line) alongside the data from all of the Upper Hunter Air Quality Monitoring Network (UHAQMN) stations. The figure shows that all of the Upper Hunter stations recorded similar trends of elevated PM_{10} levels from approximately 1:00am to 10:00am during north-westerly winds with speeds of approximately 2-5m/s. The Tamworth monitor also recorded generally similar trends however the elevated PM_{10} levels at Tamworth persisted until 3pm.

The Namoi monitors shown in **Figure 8-1** also recorded similar trends to the UHAQMN monitors however the levels were a little lower.

The available data indicate that there were elevated PM_{10} levels recorded by the monitors in the Hunter Valley and in the Namoi region, including the Tamworth monitor. However lower wind speeds, and higher PM_{10} levels recorded later in the day (11:00 to 3:00pm), occurred at Tamworth.

It is not certain what the cause of the elevated dust levels in Tamworth may have been. A plausible explanation may be that a regional elevated dust event reached the Namoi and Hunter Valley monitors at approximately the same time, i.e. arriving when the westerly winds shift to north westerly. Such an event, combined with the low wind speed at Tamworth, which is further east than the other Namoi monitors, may explain why the Tamworth monitor experienced elevated dust levels later than the other sites. However the elevated levels at Tamworth could also have been affected by localised dust sources in or near the town, or other unknown factors. $\text{PM}_{2.5}$ levels remained low through the period which indicates that the elevated PM_{10} levels were unlikely to be due to wood smoke or open fires.




Figure 8-1: Analysis of elevated PM₁₀ levels on 10 April 2017 - Tamworth

The Tamworth monitor recorded elevated PM₁₀ levels during a period of winds from the west and northwest from approximately 4am to 3pm.




Figure 8-2: UHAQMN and Tamworth monitoring data on 10 April 2017

All of the UHAQMN stations recorded very similar trends of elevated PM₁₀ levels from approximately 1:00am to 10:00am during north-westerly winds with speeds of approximately 2-5m/s.

9 CONCLUSIONS

The results indicate that the monitoring stations generally recorded very good air quality in autumn 2017.

The Tamworth TEOM monitor recorded a 24-hour average level slightly above the PM₁₀ criterion of 50µg/m³ on 10 April 2017. All 24-hour average PM_{2.5} data recorded at the Namoi monitoring sites were below the criterion of 25µg/m³.

Relative to the Air Quality Index, during autumn 2017:

- ✦ The measured PM₁₀ levels were very good 67.6% to 97.8% of the time in autumn 2017. Levels were very good or good 94.1% to 100% of the time. The Wil-gai monitor recorded fair levels on four days (5.9%), and the Maules Creek and Tamworth monitors both recorded one day (1.1%) with fair levels. The Tamworth monitor recorded one day (1.1%) with poor levels;
- ✦ The measured levels of PM_{2.5} were generally very good in autumn 2017. The Maules Creek monitor recorded very good levels 100% of the time and the other monitors recorded very good levels between 65% (Breeza) and 93% (Wil-gai) of the time, with good levels recorded in remaining periods.

On this basis it can be concluded that the air quality in the Namoi region was generally very good in autumn 2017.

10 REFERENCES

NEPC (2001)

"National Environment Protection (Ambient Air Quality) Measure Technical Paper No. 5 Data Collection and Handling", National Environment Protection Council, May 2001.

NEPC (2016)

"National Environment Protection (Ambient Air Quality) Measure", National Environment Protection Council, March 2016.

NSW EPA (2017)

"Approved Methods for the Modelling and Assessment of Air Pollutants in NSW", State of NSW and Environment Protection Authority, January 2017.

Appendix A

How to read a windrose




Figure A-1: How to read a windrose

Appendix B
Monitoring Data (Graphical)




Figure B-1: Breeza PM₁₀ (1-hour, 24-hour and rolling annual average) concentration – Autumn 2017

Rolling annual average levels are based on the previous 12-month periods of data.




Figure B-2: Maules Creek PM₁₀ (1-hour, 24-hour and rolling annual average) concentration – Autumn 2017




Figure B-3: Werris Creek PM₁₀ (1-hour, 24-hour and rolling annual average) concentration – Autumn 2017




Figure B-4: Wil-gai PM₁₀ (1-hour, 24-hour and rolling annual average) concentration – Autumn 2017




Figure B-5: Tamworth PM₁₀ (1-hour, 24-hour and rolling annual average) concentration – Autumn 2017




Figure B-6: Breeza PM_{2.5} (1-hour, 24-hour and rolling annual average) concentration – Autumn 2017




Figure B-7: Maules Creek PM_{2.5} (1-hour, 24-hour and rolling annual average) concentration – Autumn 2017




Figure B-8: Werris Creek PM_{2.5} (1-hour, 24-hour and rolling annual average) concentration – Autumn 2017




Figure B-9: Wil-gai PM_{2.5} (1-hour, 24-hour and rolling annual average) concentration – Autumn 2017




Figure B-10: Tamworth PM_{2.5} (1-hour, 24-hour and rolling annual average) concentration – Autumn 2017

Appendix C
Monitoring Data (Tabulated)

Table C-1: 24-hour average monitoring data

Date	PM ₁₀ (µg/m ³)					PM _{2.5} (µg/m ³)				
	Breeza	Maules Creek	Werris Creek	Wil-gai	Tamworth	Breeza	Maules Creek	Werris Creek	Wil-gai	Tamworth
01/03/2017	14.7	3.4	2.1	6.0	7.3	10.8	0.1	-0.4	2.8	5.0
02/03/2017	16.0	5.4	5.3	6.9	9.2	12.0	1.0	0.8	3.1	3.4
03/03/2017	17.6	8.9	4.9	7.8	9.9	13.1	3.0	2.2	3.5	4.4
04/03/2017	12.7	2.6	2.9	4.1	8.4	9.1	-0.3	1.0	1.7	3.9
05/03/2017	10.9	4.2	5.8	4.1	6.3	7.3	2.5	3.9	2.9	3.6
06/03/2017	14.6	7.7	5.3	7.0	9.2	11.0	3.4	2.5	3.2	5.1
07/03/2017	15.3	11.4	7.7	10.3	14.0	10.1	2.7	2.9	2.9	5.9
08/03/2017	14.0	8.4	5.0	8.8	9.9	10.3	1.3	1.5	2.8	5.3
09/03/2017	24.1	7.3	8.3	8.8	14.1	-	1.2	4.7	2.5	5.8
10/03/2017	-	25.6	8.7	13.6	15.5	-	4.7	3.1	4.3	6.3
11/03/2017	7.0	11.1	9.7	15.5	15.2	3.0	2.6	4.5	4.9	7.2
12/03/2017	5.8	10.9	10.4	33.3	15.5	3.1	2.5	4.8	6.4	5.6
13/03/2017	-	-	6.2	7.3	12.4	-	-	2.7	3.1	-
14/03/2017	-	0.3	1.1	2.1	7.9	-	-0.4	-0.3	1.5	4.2
15/03/2017	-	2.2	2.2	3.3	7.3	-	0.5	-0.6	1.7	-
16/03/2017	8.1	3.0	4.4	5.4	9.9	6.1	0.6	1.0	2.3	4.2
17/03/2017	5.8	10.4	6.0	8.7	12.1	5.5	4.3	3.6	3.0	3.6
18/03/2017	9.3	4.0	2.4	5.6	7.3	7.9	2.1	-0.1	3.0	-
19/03/2017	4.7	5.3	3.5	6.0	8.6	3.0	2.9	1.4	3.6	5.3
20/03/2017	6.8	3.6	4.6	-	7.8	5.6	1.8	1.9	-	5.2
21/03/2017	12.2	4.1	2.2	-	9.9	10.5	2.3	-0.1	-	1.9
22/03/2017	7.6	9.0	5.7	-	9.5	5.6	3.3	2.8	-	4.6
23/03/2017	12.7	7.5	8.8	-	10.6	10.6	2.3	4.8	-	5.1
24/03/2017	8.1	8.4	5.4	-	6.4	6.4	3.6	2.6	-	4.0
25/03/2017	5.2	21.7	4.0	-	11.5	3.0	1.6	1.1	-	3.3
26/03/2017	11.2	6.7	5.6	-	11.6	8.7	2.9	2.5	-	5.8
27/03/2017	10.5	8.9	7.7	-	16.4	8.1	4.4	3.3	-	7.1
28/03/2017	8.8	10.8	11.4	-	21.6	6.1	5.3	5.4	-	4.5
29/03/2017	12.8	11.6	11.1	-	34.0	8.6	5.3	5.1	-	6.0
30/03/2017	13.7	-	5.1	-	13.6	11.6	-	2.6	-	7.3
31/03/2017	4.5	9.3	8.3	-	17.6	1.6	2.2	4.4	-	6.5
01/04/2017	6.0	3.7	8.7	-	15.7	3.8	2.1	6.4	-	4.4
02/04/2017	9.2	4.6	7.8	-	15.1	6.4	1.3	4.0	-	5.3
03/04/2017	7.7	4.3	5.8	5.7	13.3	5.1	1.3	4.0	2.5	4.1
04/04/2017	4.4	1.5	4.7	5.9	12.9	2.4	-0.8	2.0	1.4	2.9
05/04/2017	5.8	2.9	5.5	14.8	13.5	3.8	-0.7	3.3	2.9	5.3
06/04/2017	6.3	3.1	4.1	6.3	9.7	4.5	-0.8	2.2	1.8	-
07/04/2017	6.1	1.1	6.7	7.7	11.5	4.0	-0.7	4.4	1.3	6.0
08/04/2017	6.4	3.6	9.2	7.4	8.1	3.3	0.3	6.4	0.1	3.6
09/04/2017	8.5	6.8	6.3	14.8	10.8	4.6	-1.7	3.7	3.5	2.9
10/04/2017	18.3	34.7	25.1	39.4	54.1	7.5	2.7	4.1	7.1	10.9
11/04/2017	10.7	17.5	11.5	22.4	21.1	6.2	3.3	3.9	4.8	3.2
12/04/2017	7.5	5.4	5.6	9.3	10.7	4.8	0.0	3.4	2.5	3.3
13/04/2017	9.3	9.1	4.2	8.6	12.4	6.1	1.4	0.9	3.5	5.3
14/04/2017	10.0	11.0	11.5	21.3	14.5	6.7	4.0	8.1	6.3	8.8
15/04/2017	15.4	10.5	8.1	19.3	16.3	10.0	4.2	5.8	6.8	8.3
16/04/2017	18.5	11.8	15.3	25.0	20.6	13.3	5.4	11.4	8.1	10.8
17/04/2017	20.1	13.9	13.5	28.9	24.5	13.4	5.9	8.8	10.2	11.6
18/04/2017	18.5	16.6	12.4	26.1	19.8	13.3	6.9	7.3	10.4	8.8
19/04/2017	11.6	8.8	9.6	-	17.7	8.3	2.7	5.2	-	5.5
20/04/2017	11.1	2.8	10.1	-	11.4	6.1	-0.7	5.7	-	4.3
21/04/2017	9.7	4.8	8.9	-	10.5	5.1	-0.2	4.7	-	2.7
22/04/2017	8.8	7.1	5.1	12.0	10.7	6.1	0.7	1.7	2.2	3.8
23/04/2017	10.3	9.5	8.1	17.6	15.6	6.8	1.3	4.7	4.1	5.5
24/04/2017	16.0	16.8	14.3	41.8	18.0	6.7	1.0	8.8	11.3	6.1
25/04/2017	13.0	5.9	6.3	14.6	19.0	5.3	0.2	-	4.0	1.8
26/04/2017	11.6	3.0	0.9	5.5	5.0	7.4	-0.3	-	2.2	8.4
27/04/2017	8.0	2.3	8.0	4.3	11.8	4.2	0.6	6.5	1.5	6.2
28/04/2017	8.8	3.3	8.8	-	11.0	4.3	-0.2	6.5	-	6.0
29/04/2017	12.5	4.0	10.5	-	12.8	7.5	1.1	8.3	-	8.4
30/04/2017	10.4	5.3	11.1	-	12.0	6.9	2.1	9.8	-	6.7
01/05/2017	18.5	3.8	11.8	-	15.1	8.7	-0.3	9.0	-	4.7
02/05/2017	19.6	6.9	14.1	-	19.2	9.9	1.8	8.9	-	7.2
03/05/2017	18.1	9.3	12.9	-	12.5	8.5	2.9	8.9	-	6.9

Date	PM ₁₀ (µg/m ³)					PM _{2.5} (µg/m ³)				
	Breeza	Maules Creek	Werris Creek	Wil-gai	Tamworth	Breeza	Maules Creek	Werris Creek	Wil-gai	Tamworth
04/05/2017	12.8	4.0	5.9	-	6.7	5.2	0.2	3.3	-	2.3
05/05/2017	11.6	4.9	7.7	11.2	11.7	5.4	0.9	5.4	1.1	5.4
06/05/2017	13.5	4.7	8.5	16.8	15.1	7.3	0.3	5.4	3.1	8.6
07/05/2017	11.0	7.4	12.1	15.6	17.2	5.2	1.9	8.0	4.5	9.6
08/05/2017	13.9	11.1	13.2	32.1	19.2	6.9	3.1	7.0	7.3	7.7
09/05/2017	15.7	15.0	13.2	26.3	15.2	10.4	3.6	9.2	7.3	7.6
10/05/2017	15.4	15.8	11.8	27.6	18.5	10.3	4.5	8.3	7.9	9.3
11/05/2017	18.3	-	14.1	40.6	21.4	13.3	-	9.7	9.8	10.4
12/05/2017	17.9	9.6	17.0	24.8	16.3	12.3	2.3	10.4	6.6	8.3
13/05/2017	15.3	7.1	8.0	21.3	12.7	8.8	1.2	4.0	3.8	2.6
14/05/2017	14.4	7.4	6.6	18.2	7.8	8.9	0.2	4.6	4.6	6.1
15/05/2017	15.0	7.7	10.7	22.9	17.5	10.4	3.9	8.5	7.7	10.7
16/05/2017	13.2	8.2	12.6	37.4	22.0	8.7	3.1	9.0	8.4	13.0
17/05/2017	14.2	15.5	12.7	24.4	20.9	8.3	5.2	8.2	6.9	10.2
18/05/2017	18.9	14.2	11.8	28.0	17.4	12.7	5.2	8.2	8.1	6.7
19/05/2017	8.8	1.2	4.5	4.5	8.1	5.0	0.6	2.9	2.2	2.9
20/05/2017	5.6	-2.0	3.4	4.5	7.4	-	-1.6	2.5	2.9	-
21/05/2017	14.3	3.9	6.4	4.9	12.5	11.1	2.9	5.0	2.9	9.7
22/05/2017	5.4	3.9	6.9	7.5	13.1	2.1	0.2	5.4	4.5	8.6
23/05/2017	7.0	-	7.5	7.1	10.2	6.3	-	6.2	4.4	-
24/05/2017	4.2	-	6.5	5.3	9.4	1.8	-	4.1	2.9	9.9
25/05/2017	7.4	3.5	7.1	5.8	13.6	6.3	1.5	4.7	2.2	9.3
26/05/2017	9.0	5.5	10.4	8.9	18.3	6.7	2.5	7.2	3.8	11.4
27/05/2017	10.6	5.3	10.9	12.3	18.2	7.6	2.0	8.6	6.4	13.5
28/05/2017	8.0	4.0	7.4	7.3	11.8	5.3	2.3	5.8	4.3	-
29/05/2017	7.4	2.0	6.7	5.0	13.2	5.8	0.2	5.0	1.8	10.5
30/05/2017	11.7	4.8	10.6	6.9	16.2	10.0	1.8	7.5	1.9	14.9
31/05/2017	-	3.1	8.4	7.7	13.7	-	1.2	5.7	1.8	9.8

- Not applicable